



# OROSORB

With

Trans  Phloem Guide  
technology



Let's grow greener



Rovensa  
Next



## ADVANCED ADJUVANT TECHNOLOGY

### Description

OROSORB™ is an excellent tank mix partner for your pesticide and fertilizer sprays. It has superior spreading and penetration properties compared to other adjuvants such as crop oils (COC), methylated seed oils (MSO), high surfactant methylated seed oils (HSMSO), and non-ionic surfactants (NIS). And now with TransPhloem™ technology, it ensures faster delivery of active ingredients to target sites within plants. This means OROSORB can improve the efficacy and performance of systemic pesticides and foliar nutrients sprays compared to these other types of adjuvants. Using OROSORB simplifies choosing the right adjuvant for the job and eliminates the need to have several adjuvant products on hand. It should be your adjuvant of choice in every spray tank.

### Herbicide programs

- TransPhloem technology delivers MORE post-emergent, systemic herbicide to the weeds' roots FASTER than other types of adjuvants
- Superior spreading, wetting, and penetrating properties compared to other adjuvants
- Enhanced rain fastness due to quick cuticle penetration
- Ideal for herbicide resistance management programs

### Insecticide/ Fungicide/ Miticide Programs

- Penetrates waxy cuticles to move systemic pesticides into the plant quickly
- Is an excellent spreader, at low rates, for contact pesticides
- Knocks down mite webbing and leaves no sticky residue to attract dust and cause mite are-ups
- Unlike crop oil applications, it does not decrease stomatal conductance, reduce transpiration and photosynthesis, or increase plant stress
- If use of a MSO or COC adjuvant is required by the pesticide label, OROSORB can be added to improve spreading and penetration

### Foliar Nutritional programs

- Increases nutrient uptake into the plant
- Ensures complete coverage for better nutrient distribution

### What is Transphloem technology and why is it so important?

*TransPhloem technology accelerates the movement of pesticide active ingredients and nutrients into a plant's phloem for translocation throughout the plant.* An Oro Agri commissioned study conducted by the University of Illinois validates that TransPhloem technology improves the translocation of a glyphosate application (Roundup PowerMAX®). Using a C14 radio isotope of glyphosate, the study concluded that an improved OROSORB adjuvant was absorbed into the leaf quicker, and moved a greater percentage of applied herbicide to the roots (the active site for glyphosate's mode of action) faster than the other major types of adjuvants.


TransPhloem technology, combined with the superior spreading and penetrating properties of OROSORB, can boost your systemic herbicide's performance and give you a better return on your herbicide program investment.

# How OROSORB can improve your post-emergent weed control?

## The Challenge

Delivering a killing dose of a systemic herbicide to the roots of emerged weeds is not a easy task. Many weeds have hairy or waxy surfaces, like this fat hen leaf, that provide a repellent barrier for post-emergent herbicide applications. Without the right adjuvant in your tank mix a significant portion of your herbicide spray could be lost to run-off or to rain.


## Fat Hen Leaf Waxy Surface



Univ. Arizona – Dr. William McCloskey

Additionally, the herbicide and adjuvant need to translocate to the roots via the phloem, part of the plant's vascular system. Within the phloem are cells that regulate and can restrict the movement of water, nutrients and materials, including herbicides. Moving a herbicide solution into the phloem and then having it translocate effectively to the roots, through the phloem, is another challenge to achieving excellent weed control.

## Phloem within a plants vascular system






## Movement of Materials within the Phloem

Phloem allows materials to flow both up and down the plants vascular system.

A lateral cut of a stem illustrates the phloem and xylem vascular bundles.

The phloem carries materials throughout the plant including down to the root system.



Univ. Arizona – Dr. William McCloskey

## The Solution

The critical first step in post-emergent weed control is getting good coverage of the leaf surface with your herbicide spray. OROSORB, because of its superior surface tension reduction property, provides excellent wetting and coverage of the leaf. It also delivers a greater number of droplets and a more uniform droplet size compared to other adjuvant types that contributes to its excellent spray coverage.

### 1. Superior Spreading



**OROSORB**



**Untreated**



**Droplet Study – Per 2 cm<sup>2</sup>**



OROSORB


HSMSO

NIS 90

COC


## 2. Superior Penetration

Leaf surfaces often present a formidable barrier to the delivery of systemic herbicides into the plant, as illustrated by the magnified photo of the fat hen leaf on pg. 2. However, OROSORB overcomes these barriers with a novel formulation that allows movement through the epicuticular wax on the leaf surface and between cells inside the leaf.



## 3. Superior Absorption

In the Oro Agri-commissioned study, OROSORB, with TransPhloem technology, was shown to move a Roundup PowerMAX solution into the leaf more than twice as fast as a Roundup PowerMAX + HSMSO solution or Roundup PowerMAX alone. In as little as 3 minutes the OROSORB solution is thoroughly absorbed into the leaf, delivering more herbicide for quick translocation and providing excellent rainfastness.



HSMSO 0.25% + Roundup PowerMAX  
Roundup PowerMAX w/o adjuvant


HSMSO 0.25% + Roundup PowerMAX  
Roundup PowerMAX w/o adjuvant  
OROSORB + Roundup PowerMAX

HSMSO 0.25% + Roundup PowerMAX  
Roundup PowerMAX w/o adjuvant  
OROSORB + Roundup PowerMAX

### 3. Superior Absorption Continued

The result of the excellent leaf absorption and penetration of OROSORB is shown in this photo from the recent study. A radio isotope solution of glyphosate, with and without OROSORB, was applied to a fat hen leaf. After only 15 minutes, the OROSORB application had penetrated, and was distributed, completely through the leaf whereas the penetration of the glyphosate alone was spotty. Also very significant is the movement of the OROSORB solution into the petiole for translocation only 15 minutes after application.


#### Radio isotope herbicide absorption – Fat Hen: 15 Minutes after application



### 4. Superior Translocation

TransPhloem technology effects the herbicide and adjuvant movement when it moves into the main stem of the weed. Research shows that when the improved OROSORB is added to glyphosate a higher percentage of applied herbicide is translocated, more quickly, compared to glyphosate alone or mixed with other adjuvants. As importantly, OROSORB delivered more herbicide to the critical lower parts of the plant, and less upward in the plant, than any other adjuvant.

#### Translocation of glyphosate into fat hen plan sections at 24 Hours






Further proof of the superior translocation provided by TransPhloem technology is shown by these phosphor images of the C14 radio isotope glyphosate concentration in fat hen plants. A single leaf was treated with the glyphosate/adjutant solutions. At 12 hours, the plants were uprooted and the phosphor images were taken. Only those parts of the plant that have some concentration of C14 glyphosate show up in the image. Red areas indicate high concentrations, green areas show medium concentrations, and blue areas show low concentrations. The OROSORB-treated plant has more area visible signifying the C14 solution has moved into more parts of the plant than other adjuvants. Also, the OROSORB solution has the most green areas and very little red areas on the treated leaf, indicating the glyphosate has moved out of the leaf into other areas of the plant. The OROSORB plant also has the largest amount of red area in the roots, showing a high concentration of glyphosate has been translocated to the roots in 12 hours.


### **FAT Hen: 12 Hr after application – Radio isotope phosphor images**



**OROSORB**




**MSO**




**HSMSO**



**GLYPHOSATE**




**COC**




**NIS 90**

## OROSORB effect on Scald infection in Barley

Scald infection at 31DAAB



Addition of OROSORB to Radial significantly reduced Scald infection.




Trial Location : Cressy, Tasmania  
Sowing date: 05/05/21  
Spray volume: 100 L/ha  
6 Replications

Crop: Barley, cv. Planet  
Pest: Net blotch and Scald  
Pest pressure: moderate  
Grower standard: Radial


Application A: 05/08/21, 12°C, 82%RH  
Application B: 06/09/21, 17°C, 44%RH

## OROSORB effect on Net Blotch infection in Barley

Net blotch infection at 31DAAB



Addition of OROSORB to Radial significantly reduced net blotch infection.



Trial Location : Cressy, Tasmania  
Sowing date: 05/05/21  
Spray volume: 100 L/ha  
6 Replications

Crop: Barley, cv. Planet  
Pest: Net blotch and Scald  
Pest pressure: moderate  
Grower standard: Radial


Application A: 05/08/21, 12°C, 82%RH  
Application B: 06/09/21, 17°C, 44%RH



# Effect of various adjuvants on control of Fleabane

% Control with glyphosate + 2,4-D Amine + trichlopyr

Control on fleabane



- Addition of OROSORB at 0.2% v/v is equivalent to HSMOC for the control of fleabane.
- Herbicide mixture with Orosorb achieve quicker control at 49DAA when compared to HSMOC.




All treatments with full rate herbicide mixture:  
Weedmaster DST (47% glyphosate) @ 291 g a.i./ac  
2,4-D amine (70% ) @ 189 g a.i./ac  
Garlon: (60% Trichlopyr) @ 20 g a.i./ac  
No stats done as only 3 replicates

Application date: May 29, 2020  
Water vol: 10 gal/ac (100 L/ha)

Fleabane (*Conyza bonariensis*)  
Potted weeds (5 plants/pot)

## Effect of adjuvants (OROSORB) on control of Rye grass

Ryegrass control



Weed: Rye grass (*Lolium rigidum*)  
Vetch (*Vicia sativa*)

Roundup CT applied at low rate (1L/ha) for adjuvant to perform  
Crop: Fallow

Application date: 23/10/2020  
Water vol: 97 L/ha  
4 replications

## The Results

OROSORB, with TransPhloem technology, moves systemic herbicides into the plants roots quicker and at higher concentrations than other types of adjuvants. The end result is a faster kill of weeds that rob your crop of moisture and nutrients. As shown in this controlled-environment pot study the OROSORB + glyphosate treated fat hen was killed quicker than any other adjuvant / glyphosate combination

Transphloem technology combined with superior spreading, penetration and absorption results in a faster kill.

### Burn down herbicide trial – Fat Hen: OROSORB, COC, MSO, HSMSO & NIS 90



OROSORB

COC

MSO

HSMSO

NIS 90

## Application and Usage Directions

A real benefit to growers is the versatility OROSORB adjuvant brings to spray programs. OROSORB has the capability to act as a wetter, spreader, penetrant and a translocation aid depending on the use rate. Regardless of whether you are spraying a contact or systemic pesticide or foliar nutrient OROSORB can deliver it to the desired action site — giving the active ingredients the best chance for peak performance. This makes selecting the right adjuvant simple and means you don't have to buy several adjuvants for your different spray programs.

### LABEL USE RATES

HERBICIDES, DEFOLIANTS, DESICCANTS: 100 - 250 ml (0.1% - 0.25%) per 100 litres of water.

TO CONTROL HARD TO KILL WEEDS: 400 - 800 ml (0.4 - 0.8%) per 100 litres of water.

### MIXING AND APPLICATION INSTRUCTIONS:

- Follow the mixing and application instructions for the pesticide(s) or foliar nutrient(s) being tank mixed with OROSORB.
- With constant agitation, add OROSORB to the spray tank after all other materials are thoroughly dissolved or mixed.
- OROSORB will not affect the pH of the spray solution.
- Be aware OROSORB will dissolve any residues in the tank, lines and nozzles from previous spray materials.

If these residues could cause harm to the crop being sprayed thoroughly clean your spray system prior to mixing OROSORB in the spray tank. Use a commercial tank cleaner or mix 600 ml of OROSORB in 75 litres of water and thoroughly circulate through your sprayer. Dispose of the spray solution properly.

*Always read and follow the product labels for more complete information and application directions.*



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# OROSORB

SPRAY ADJUVANT

A WETTING AGENT FOR USE WITH INSECTICIDES, MITICIDES, FUNGICIDES, HERBICIDES, PLANT GROWTH REGULATORS AND NUTRIENTS ON NON-FRUIT CROPS

ACTIVE INGREDIENT  
190.9 g/L ANIONIC SURFACTANTS  
81.5 g/L ALCOHOL ETHOXYSULFATE

In a transport emergency, dial 900 police or fire brigade.

For hazardous materials or dangerous goods incidents ONLY  
(spill, leak, fire, exposure, or accident), call Chemtrec at +61(0)12 9037 2964.

SITUATION	RATE PER 100 L	CRITICAL COMMENTS
Insecticides, Fungicides, Miticides	50 - 200 mL (0.05% - 0.2%)	Use the higher rate on hairy, waxy, dusty or hard to wet surfaces. Prior to using the product read all directions on this label and on the label of the agricultural chemical being used, including the Safety Directions, First Aid Instructions, Precaution statements and any Warning statements.
Plant growth regulators and foliar applied nutrients	25-100 mL (0.025% - 0.1%)	
Herbicides	100-250 mL (0.1% - 0.25%)	
Hard to kill weeds	400-800 mL (0.4% - 0.8%)	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL, UNLESS AUTHORIZED UNDER APPLICABLE LEGISLATION.

**APÉPÉ FREE**

NET CONTENTS 10 L

MANUFACTURED BY  
21st Century, Inc.  
2780 S. Main Ave., Fresno, CA 93726 USA  
21st Century, Inc. All Rights Reserved  
Copyright © Nov. 2022 02029117  
Lot number: See packaging

**Rovensa Next**

DISTRIBUTED BY  
ROVENSA INC PTY LTD  
Level 10, 51 Chisholm Street  
Melbourne VIC 3207 Australia  
Ph: +61(0)3 556 350

Phone: 1300 595 000  
[Info.au@rovensanext.com](mailto:Info.au@rovensanext.com)  
<https://www.rovensanext-au-nz.com/>

